AUTHOR INDEX

Abe, M., Ito, S.-i., Kimoto, M., Hayashi, R. and Nishimune, T.
Molecular studies on thiaminase I (909) 213

Adolph, K.W.
ADPribosylation of nuclear proteins labeled with [^{3}H]adenosine: changes during the HeLa cell cycle (909) 222

Anderer, F.A., see Thomas, L. (909) 173

Andrews, G.K., Varma, S. and Ebner, K.E.
Regulation of expression of c-*fos* and c-*myc* in rat lymphoma Nb-2 cells (909) 231

Angal, S., see Bodmer, M.W. (909) 237

Baer, G., see Heumann, H. (909) 126

Bauer, P.I., see McLick, J. (909) 71

Bernués, J., see Espel, E. (909) 190

Bianchi, M.S., see Bianchi, N.O. (909) 245

Bianchi, N.O., Peltomäki, P., Bianchi, M.S., Knuutila, S. and De la Chapelle, A.
The pattern of methylation in rearranged and germ-line human immunoglobulin constant μ genes (909) 245

Bodmer, M.W., Angal, S., Yarranton, G.T., Harris, T.J.R., Lyons, A., King, D.J., Pieroni, G., Riviere, C., Verger, R. and Lowe, P.A.
Molecular cloning of a human gastric lipase and expression of the enzyme in yeast (909) 237

Buckingham, K., Chung, D.-G., Neilson, T. and Ganoza, M.C.
Recognition of translational termination signals (909) 92

Butkus, V., Klimašauskas, S., Petrauskienė, L., Maneliené, Z., Lebionka, A. and Janulaitis, A.
Interaction of *Alu*I, *Cfr*6I and *Pvu*II restriction-modification enzymes with substrates containing either N^4-methylcytosine or 5-methylcytosine (909) 201

Cassidy, B., Haglund, R. and Rothblum, L.I.
Regions upstream from the core promoter of the rat ribosomal gene are required for the formation of a stable transcription initiation complex by RNA polymerase I in vitro (909) 133

Chang, D.D., Fisher, R.P. and Clayton, D.A.
Roles for a promoter and RNA processing in the synthesis of mitochondrial displacement-loop strands (909) 85

Chung, D.-G., see Buckingham, K. (909) 92

Clayton, D.A., see Chang, D D. (909) 85

Davie, J.R. and Nickel, B.E.
The ubiquitinated histone species are enriched in histone H1-depleted chromatin regions (909) 183

De la Chapelle, A., see Bianchi, N.O. (909) 245

Ebner, K.E., see Andrews, G.K. (909) 231

Endo, H., see Kohnoe, S. (909) 107

Espel, E., Bernués, J., Guasch, M.D., Querol, E., Plana, M. and Itarte, E.
Phosphorylation of high-mobility-group protein 14 by two specific kinases modifies its interaction with histone oligomers in free solution (909) 190

Fisher, R.P., see Chang, D.D. (909) 85

Flavell, D., see Julien, J.-P. (909) 10

Fox, K.R. and Waring, M.J.
The use of micrococcal nuclease as a probe for drug-binding sites on DNA (909) 145

Fujiki, Y., see Rachubinski, R.A. (909) 35

Ganoza, M.C., see Buckingham, K. (909) 92

Glusker, J.P., see McLick, J. (909) 71

Grosveld, F., see Julien, J.-P. (909) 10

Guasch, M.D., see Espel, E. (909) 190

Haglund, R., see Cassidy, B. (909) 133

Hakam, A., see McLick, J. (909) 71

Harano, T., see Liu, J.-Z. (909) 208

Harris, T.J.R., see Bodmer, M.W. (909) 237

Hayashi, R., see Abe, M. (909) 213

Heumann, H., Lederer, H., Kammerer, W., Palm, P., Metzger, W. and Baer, G.
Large-scale preparation of a DNA fragment containing the strong promoter A1 of the phage T7 (909) 126

Ho, K.-C. and Stafford, D.W.
Nucleotide sequence of two 3′-termini of rRNA in the sea urchin *Lytechinus variegatus* (909) 156

Houssier, C., see Marquet, R. (909) 165

Huisman, T.H.J., see Liu, J.-Z. (909) 208

Ide, T., see Takasuka, T. (909) 161

Ishibashi, S., see Takasuka, T. (909) 161

Ishikawa, K., see Kuzumaki, T. (909) 99

Itarte, E., see Espel, E. (909) 190

Ito, S.-i., see Abe, M. (909) 213

Janulaitis, A., see Butkus, V. (909) 201

Jardetzky, O., see Lane, A.N. (909) 58

Julien, J.-P., Grosveld, F., Yazdanbaksh, K., Flavell, D., Meijer, D. and Mushynski, W.
The structure of a human neurofilament gene (NF-L): a unique exon-intron organization in the intermediate filament gene family (909) 10

Kammerer, W., see Heumann, H. (909) 126
Karawya, E.M. and Martin, R.G.
Monkey (CV-1) mitochondrial DNA contains a unique triplication of 108 bp in the origin region (909) 30
Kimoto, M., see Abe, M. (909) 213
King, D.J., see Bodmer, M.W. (909) 237
Klimašauskas, S., see Butkus, V. (909) 201
Knuutila, S., see Bianchi, N.O. (909) 245
Kohnoe, S., Maehara, Y. and Endo, H.
A systematic survey of repetitive sequences abundantly expressed in rat tumors (909) 107
Kun, E., see McLick, J. (909) 71
Kuzumaki, T., Tanaka, T., Ishikawa, K. and Ogata, K.
Rat ribosomal protein L35a multigene family: molecular structure and characterization of three L35a-related pseudogenes (909) 99

Lanclos, K.D., see Liu, J.-Z. (909) 208
Lane, A.N., Lefèvre, J.-F. and Jardetzky, O.
The interaction of the trp repressor from *Escherichia coli* with the trp operator (909) 58
Lazarow, P.B., see Rachubinski, R.A. (909) 35
Lebionka, A., see Butkus, V. (909) 201
Lederer, H., see Heumann, H. (909) 126
Lefèvre, J.-F., see Lane, A.N. (909) 58
Linnala-Kankkunen, A., see Palvimo, J. (909) 21
Liu, J.-Z., Harano, T., Lanclos, K.D. and Huisman, T.H.J.
The β–δ crossover leading to the $\beta\delta$ hybrid gene of hemoglobin P-Nilotic is located within 54 base-pairs of the 5′ end of exon 2 or between codons 31 and 50 (909) 208
Lowe, P.A., see Bodmer, M.W. (909) 237
Lyons, A., see Bodmer, M.W. (909) 237

McLick, J., Hakam, A., Bauer, P.I., Kun, E., Zacharias, D.E. and Glusker, J.P.
Benzamide–DNA interactions: deductions from binding, enzyme kinetics and from X-ray structural analysis of a 9-ethyladenine-benzamide adduct (909) 71
Maehara, Y., see Kohnoe, S. (909) 107
Mäenpää, P.H., see Palvimo, J. (909) 21
Malcolm, A.D.B., see Woody, A.-Y.M. (909) 115
Manelienė, Z., see Butkus, V. (909) 201
Marquet, R., Wyart, A. and Houssier, C.
Influence of DNA length on spermine-induced condensation. Importance of the bending and stiffening of DNA (909) 165
Martin, R.G., see Karawya, E.M. (909) 30
Meijer, D., see Julien, J.-P. (909) 10
Metzger, W., see Heumann, H. (909) 126
Mushynski, W., see Julien, J.-P. (909) 10

Neilson, T., see Buckingham, K. (909) 92
Nickel, B.E., see Davie, J.R. (909) 183
Nishimune, T., see Abe, M. (909) 213
Nover, L., see Scharf, K.-D. (909) 44

Ogata, K., see Kuzumaki, T. (909) 99

Palm, P., see Heumann, H. (909) 126
Palvimo, J., Pohjanpelto, P., Linnala-Kankkunen, A. and Mäenpää, P.H.
Alterations in amounts and covalent modifications of low-molecular-weight chromosomal proteins in Chinese hamster ovary cells during polyamine depletion (909) 21
Peltomäki, P., see Bianchi, N.O. (909) 245
Petrauskienė, L., see Butkus, V. (909) 201
Pfeifle, J., see Thomas, L. (909) 173
Pieroni, G., see Bodmer, M.W. (909) 237
Plana, M., see Espel, E. (909) 190
Pohjanpelto, P., see Palvimo, J. (909) 21

Querol, E., see Espel, E. (909) 190

Rachubinski, R.A., Fujiki, Y. and Lazarow, P.B.
Isolation of cDNA clones coding for peroxisomal proteins of *Candida tropicalis*: identification and sequence of a clone for catalase (909) 35
Riviere, C., see Bodmer, M.W. (909) 237
Rothblum, L.I., see Cassidy, B. (909) 133

Scharf, K.-D. and Nover, L.
Control of ribosome biosynthesis in plant cell cultures under heat shock conditions. II. Ribosomal proteins (909) 44
Stafford, D.W., see Ho, K.-C. (909) 156

Takasuka, T., Ishibashi, S. and Ide, T.
Expression of cell-cycle-dependent genes in serum stimulated cells whose entry into S phase is blocked by cytochalasin D (909) 161
Tanaka, T., see Kuzumaki, T. (909) 99
Thomas, L., Pfeifle, J. and Anderer, F.A.
Human SS-B/LA autoantigen contains a covalent protein–RNA linkage (909) 173

Varma, S., see Andrews, G.K. (909) 231
Verger, R., see Bodmer, M.W. (909) 237

Wang, J.C.
Recent studies of DNA topoisomerases (909) 1
Waring, M.J., see Fox, K.R. (909) 145
Woody, A.-Y.M., Woody, R.W. and Malcolm, A.D.B.
Effects of ppGpp on transcription by DNA-dependent RNA polymerase from *Escherichia coli:* circular dichroism, absorption and specific transcription studies (909) 115
Woody, R.W., see Woody, A.-Y.M. (909) 115
Wyart, A., see Marquet, R. (909) 165

Yarranton, G.T., see Bodmer, M.W. (909) 237
Yazdanbaksh, K., see Julien, J.-P. (909) 10

Zacharias, D.E., see McLick, J. (909) 71